

(19) 世界知的所有権機関
国際事務局(43) 国際公開日
2001 年 10 月 11 日 (11.10.2001)

PCT

(10) 国際公開番号
WO 01/74774 A1

(51) 国際特許分類: C07D 209/12, 209/42, 401/12, 401/14, 409/12, 417/12, 417/14, 471/04, 487/04, 495/04, 498/04, 513/04, 513/14, 519/00, A61K 31/404, 31/428, 31/429, 31/4365, 31/437, 31/4439, 31/454, 31/4545, 31/4725, 31/5025, 31/5377, A61P 7/02, 9/10

LTD.) [JP/JP]; 〒103-8234 東京都中央区日本橋3丁目14番10号 Tokyo (JP).

(21) 国際出願番号: PCT/JP01/02945

(22) 国際出願日: 2001 年 4 月 5 日 (05.04.2001)

(25) 国際出願の言語: 日本語

(26) 国際公開の言語: 日本語

(30) 優先権データ:
特願2000-108047 2000 年 4 月 5 日 (05.04.2000) JP

(72) 発明者; および

(75) 発明者/出願人 (米国についてのみ): 吉野利治 (YOSHINO, Toshiharu) [JP/JP]. 永田 勉 (NAGATA, Tsutomu) [JP/JP]. 萩野谷憲康 (HAGINOYA, Noriyasu) [JP/JP]. 吉川謙次 (YOSHIKAWA, Kenji) [JP/JP]. 菅野英幸 (KANNO, Hideyuki) [JP/JP]. 永持雅敏 (NAGAMOCHI, Masatoshi) [JP/JP]; 〒134-8630 東京都江戸川区北葛西1丁目16-13 第一製薬株式会社 東京研究開発センター内 Tokyo (JP).

(74) 代理人: 有賀三幸, 外 (ARUGA, Mitsuyuki et al.); 〒103-0013 東京都中央区日本橋人形町1丁目3番6号 共同ビル Tokyo (JP).


(71) 出願人 (米国を除く全ての指定国について): 第一製薬株式会社 (DAIICHI PHARMACEUTICAL CO.,

(81) 指定国 (国内): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK,

[続葉有]

(54) Title: ETHYLENEDIAMINE DERIVATIVES

(54) 発明の名称: エチレンジアミン誘導体





(2)

(57) Abstract: Compounds of the general formula (1): $Q^1-Q^2-C(=O)-N(R^1)-Q^3-N(R^2)-T^1-Q^4$; and drugs which contain the compounds and are efficacious for thrombosis and embolism. In said formula, wherein R^1 and R^2 are each H or the like; Q^1 is an aromatic ring, a heterocycle, or the like; Q^2 is a single bond, an aromatic ring, a heterocycle, or the like; Q^3 is a group of the general formula (2), or the like; Q^4 is an aromatic ring, a heterocycle, or the like; and T^1 is $-CO-$ or $-SO_2-$.

(57) 要約:

本発明は式 (1)

(式中、 R^1 および R^2 はHなど; Q^1 は芳香環、複素環など; Q^2 は単結合、芳香環、複素環など; Q^3 はなど; Q^4 は芳香環、複素環など; T^1 は $-CO-$ または $-SO_2-$ を示す)

で表される化合物およびこれを含有する血栓症、塞栓症に有効な医薬に関する。

Derwent WPI records



FILE 'HCA' ENTERED AT 17:50:52 ON 21 MAY 2004
E WO2001074774/PN

L3 1 S E27
L4 3 S WO2003000657/PN
L5 3 S WO2003000680/PN
L6 4 S L3-L5

FILE 'WPIX' ENTERED AT 17:53:21 ON 21 MAY 2004

L7 3 S L6

L7 ANSWER 1 OF 3 WPIX COPYRIGHT 2004 THOMSON DERWENT on STN
AN 2003-256264 [25] WPIX Full-text
CR 2003-247888 [24]; 2003-312743 [30]
TI New diamine compounds are activated blood factor X inhibitors used for
treating e.g. cerebral or myocardial infarction and angina.
PA (DAUC) DAIICHI PHARM CO LTD
PI WO 2003000657 A1 20030103 (200325)* JA 788 C07D209-42 <--
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ
NL OA PT SD SE SL SZ TR TZ UG ZM ZW
W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK
DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR
KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT
RO RU SD SE SG SI SK SL TJ TM TN TR TT TZ UA UG US UZ VN YU ZA ZM
ZW
PRAI JP 2001-398708 20011228; JP 2001-187105 20010620;
JP 2001-243046 20010809; JP 2001-311808 20011009
TECH UPTX: 20030410



TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preparation: (I) Are prepared e.g.
by reacting an amine of formula (IV) with an acid of formula (V).

AB WO2003000657 A UPAB: 20030513

NOVELTY - Diamine compounds (I) are new.

DETAILED DESCRIPTION - Diamine compounds of formula (I) and their salts, solvates and N-oxides are new. R1, R2 = H, OH, alkyl or O-alkyl; Q1 = 5-6C carbocyclyl, 5-7 membered heterocyclyl or bi- or tri-cyclic carbocyclyl or heterocyclyl (all optionally unsaturated and optionally substituted);

Q2 = a bond, or divalent 5-6C carbocyclyl, 5-7 membered heterocyclyl or bi- or tricyclic carbocyclyl or heterocyclyl (all optionally unsaturated and optionally substituted); Q3 = a group of formula (i);

Q5 = 1-8C alkylene, 2-8C alkenylene or (CH2)mCH2ACH2(CH2)n; m, n = 0-3;

A = O, N, S, SO, SO2, NH, ONH, NHH, SNH, SONH or SO2NH; R3, R4 = H, OH, alkyl (optionally substituted by CN, NH2, NHA, NAA, acyl, acylamino, O-alkyl, OH, COOH, COO-alkyl, NHCOOA, N-alkenylcarbamoyl, N-alkenyl-N-alkyl-carbamoyl, CONHO-alkyl, CON(A)OA, SO2A, CONH2, OCONH2, OCONHA, OCONAA, OCOHet, COHet, Ar1, NHSO2alkyl, NHSO2Ar, CONHSO2alkyl, CONHSO2Ar or optionally

substituted CONHA or CONAA), alkenyl, alkynyl, halo, acyl, optionally substituted acylamino, O-alkyl, COOH, COO-alkyl, NHACOOA, NHACOOH, CONH2, CONHA, CONAA, N-alkenylcarbamoyl, N-alkenyl-N-alkyl-carbamoyl, CONHO-alkyl, CON(A)OA, CON(A)3, SO2A, COHet, Ar1, NHSO2alkyl, NHSO2Ar, CONHSO2alkyl, CONHSO2Ar, oxo, OCONH2, aralkoxy, O-alkyl-COOH, acyloxy, SO2Ar, SO2-alkyl-COO-alkyl, SO2-alkyl-COOH, acyl (substituted by COO-alkyl, OH, O-alkyl, halo, COOH, NH2, acyloxy, NH-alkyl, N(alkyl)2, SO2alkyl or optionally substituted CON(alkyl)2, COO-alkyl-O-alkyl or acyloxyalkylsulfonyl, SO2-alkyl-OH, SO2O-alkyl, SO2Het or optionally substituted SO2CON(alkyl)2, or R3 + R4 = 1-5C alkylene, 2-5C alkenylene, 1-5C alkyleneoxy or carbonyldioxy; Het = optionally substituted 3-6C heterocyclyl; Ar1 = aryl or heteroaryl;

Ar = aryl;

Q4 = Ar1, arylalkenyl, arylalkynyl, heteroarylalkenyl or optionally unsaturated bi- or tricyclic carbocyclyl or heterocyclyl (all optionally substituted);

T = CO or CS;

T1 = CO, SO2, COCONR11, CSCONR11, COCSNR11, CSCSNR11, COA1NR12, CONH, CSNH, CONHNH, COA2CO, COA3CONH, COC(=NORa)NRb, CS(=NORa)NRbCON=N, CSN=N or CS;

R11, R12, Rb = H, OH, alkyl or O-alkyl; A1 = optionally substituted 1-5C alkylene; A2 = bond or A1, and

Ra = H, alkyl or alkanoyl.

ACTIVITY - Cerebroprotective; Cardiant; Antianginal; Respiratory; Anticoagulant; Thrombolytic; Vasotropic; Antiinflammatory.

MECHANISM OF ACTION - Factor X Inhibitor. In tests on rats, administration of N1-(4-chlorophenyl)-N2-((1S,2R,4S)-4-((dimethylamino)carbonyl)-2-((5-methyl-4,5,6,7-tetrahydrothiazolo(5,4-c)pyridin-2-yl)carbonyl)amino)cyclohexyl)ethanediamide (Ia) (10 mg/kg orally) reduced blood FXa levels over 4 hours by 62-96%.

USE - As activated blood factor X inhibitors for treating and preventing e.g. cerebral infarction, cerebral embolism, myocardial infarction, angina, pulmonary, embolism, Buerger's disease, deep vein thrombosis, disseminated intravascular coagulation, thrombosis following artificial flap/hip replacement or during external circulation, reocclusion, systemic inflammatory reaction syndrome, multiple organ failure or blood coagulation during blood collection. Dwg.0/0

ABEX

UPTX: 20030410

ADMINISTRATION - Administration of (I) is 0.1-200 (preferably 0.5-100) mg/kg/day orally or by injection.

EXAMPLE - 1-Hydroxybenzotriazole monohydrate (71 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloric acid (100 mg) were added to N-((1Rasterisk,2Sasterisk)-2-aminocyclopropyl)-5-chloroindole-2-carboxamide (108 mg) and lithium 5-methyl-4,5,6,7-tetrahydrothiazolo(5,4-c)pyridine-2-carboxylate (124 mg) in dimethylformamide (3 ml) and the mixture was stirred at room temperature for 8 hours. Work-up including silica gel chromatography (methylene chloride:methanol 10:1) gave 72 mg of N-((1Rasterisk,2Sasterisk)-2-((5-chloroindol-2-yl)carbonyl)amino)cyclopropyl)-5-methyl-4,5,6,7-tetrahydrothiazolo(5,4-c)pyridine-2-carboxamide hydrochloride.